

**THE HILINSKI CLINIC
OF FACIAL PLASTIC SURGERY**

John M. Hilinski, M.D.
3720 Fourth Avenue
San Diego, CA 92103
(619) 296-3223 Office
(619) 296-3224 Fax

RESTRICTED MEDICATIONS

Any surgery carries a risk of bleeding or infection in the post-operative period. The risk of bleeding can be reduced by NOT consuming medication that can affect your blood clotting capability. Please refrain from using any medication on the restricted list below for at least 14 days prior to any surgery. Although this is NOT a guarantee, avoidance of the following medications can significantly reduce chances of bleeding from and following surgery. *Tylenol is permissible.*

Advil	Empirin compound	Aloe Vera caps
Aleve	Equagesic	Bilberry
Alka Seltzer	Equagesic M	Cayenne
Alka Seltzer Plus	Excedrin	Echinacea
Anacin	Excedrin PM	5 PC-SPEC
Anacin Maximum	Fizri powder	Garlic
Anexsia with codeine	Fiorinal	Ginger
Anodynos DHC	Fiorinal with codeine	Gingko Biloba
APC	4-way cold tablets	Licorice root
Arthritis Pain formula	Gemnisyn	St. John's Wart
Arthritis Bufferin	Ibuprofen	Yohimbe
Aspirin with codeine	Leech Extract	Vitamin E
Aspirin	Measurin	
Aspirin enseals	Meprogesic	
Ascriptin	Midol	
Ascriptin A/D	Momentum	
Aspergum	Motrin	
Asphal G	Nilain	
B-A	Norgesic	
Bayer	Novhistine with APC	
Bayer decongestant	Pabirin	
Bayer timed release	PAC	
BC tablets	Panodynes analgesic	
Bufferin	Percodan	
Buff-A compound #3	Percodan demi	
CAMA	Persistin	
Capron capsules	Quiet world tablets	
COPE	Relafen	
Cogespirin	Robaxisal	
Coricidin D	Stanback tablets	
Damason P	Stanco	
Darvon N with ASA	St. Joseph Aspirin	
Darvon compound	Super Anahist	
Darvon compound 65	Synalogos DC	
Dolcin	Synalogos	
Dolene compound 65	Talwin compound	
Dolor	Traminicin	
Duradyne	Trigesic	
Duragesic	Vanquish	
Ecotrin		
Empirin with codeine		