

RESTRICTED MEDICATIONS

Any surgery carries a risk of bleeding or infection in the post-operative period. The risk of bleeding can be reduced by **not** consuming medication that can affect your blood clotting capability. Please refrain from using any medication on the restricted list for at least fourteen (14) days prior to any surgery. Although this is not a guarantee, avoidance of the medications can significantly reduce chances of bleeding from and following surgery. **Tylenol is permissible.**

Advil	Darvon compound	PAC
Aleve	Darvon compound 65	Panodynes analgesic
Alka Seltzer	Dolcin	Percodan
Alka Seltzer Plus	Dolene compound 65	Percodan demi
Anacin	Dolor	Persistin
Anacin Maximum	Duradyne	Quiet world tablets
Anexsia with codeine	Duragesic	Relafen
Anodynos DHC	Ecotrin	Robaxisal
APC	Empirin with codeine	Stanback tablets
Arthritis Pain formula	Empirin compound	Stanco
Arthritis Bufferin	Equagesic	St. Joseph Aspirin
Aspirin with codeine	Equagesic M	Super Anahist
Aspirin	Excedrin	Synalogos DC
Aspirin enseals	Excedrin PM	Synalogos
Ascriptin	Fizri powder	Talwin compound
Ascriptin A/D	Fiorinal	Traminicin
Aspergum	Fiorinal with codeine	Trigesic
Asphal G	Fish oil	Vanquish
B-A	4-way cold tablets	Viromed
Bayer	Gemisyn	Zactirin
Bayer decongestant	Ibuprofen	Zactirin compound
Bayer timed release	Indocin	
BC tablets	Leech Extract	
Bufferin	Measurin	
Buff-A compound #3	Meprogesic	
CAMA	Midol	
Capron capsules	Momentum	
COPE	Motrin	
Cogespirin	Nilain	
Coricidin D	Norgesic	
Damason P	Novhistine with APC	
Darvon N with ASA	Pabirin	

Homeopathic Medications and Supplements

Refrain from ALL homeopathic medications and supplements (*this includes any and all dietary and/or workout*

supplements).

The following are included:

Aloe Vera caps
Bilberry
Cayenne
Echinacea
5 PC-SPEC
Garlic
Ginger
Gingko Biloba
Licorice root
St. John's Wart
Yohimbe
Vitamin E

QUESTIONS?

Call our offices immediately at (619) 296-3223 if you have any questions or concerns.