

Certified American Board of Facial Plastic Surgery Certified American Board of Otolaryngology Head and Neck Surgery Harvard Medical School Graduate

RESTRICTED MEDICATIONS

Any surgery carries a risk of bleeding during or after the procedure. The risk of bleeding can be reduced by avoiding medication that can affect your blood clotting capability. Please refrain from using any medication on the restricted list for at least fourteen (14) days prior to and seven (7) days after your surgery. Although this is not a guarantee, avoidance of the medications can significantly reduce chances of unwanted complications. Tylenol is permissible within this time frame.

Advil Damason P Nilain
Aleve Darvon N with ASA Norgesic

Alka Seltzer Darvon compound Novhistine with APC

Alka Seltzer Plus Darvon compound 65 Pabirin
Anacin Dolcin PAC

Anacin Maximum Dolene compound 65 Panodynes analgesic

Anexsia with codeine Dolor Percodan
Anodynos DHC Duradyne Percodan demi
APC Duragesic Persistin

Arthritis Pain formula Ecotrin Quiet world tablets
Arthritis Bufferin Empirin with codeine Relafen

Aspirin with codeine Empirin compound Robaxisal

Aspirin Equagesic Stanback tablets

Aspirin ensealsEquagesic MStancoAscriptinExcedrinSt. Joseph AspirinAscriptin A/DExcedrin PMSuper AnahistAspergumFizri powderSynalogos DC

Asphal G Fiorinal Synalogos

B-A Fiorinal with codeine Talwin compound

Bayer Fish oil Traminicin
Bayer decongestant 4-way cold tablets Trigesic
Bayer timed release Gemnisyn Vanquish

BufferinIndocinZactirinBuff-A compound #3Leech ExtractZactirin compound

Viromed

Ibuprofen

CAMA Measurin
Capron capsules Meprogesic
COPE Midol
Cogespirin Momentum
Coricidin D Motrin

Homeopathic Medications and Supplements

BC tablets

Refrain from ALL homeopathic medications and supplements, which includes dietary and/or workout supplements. This includes, but is not limited to the following:

Aloe Vera caps 5 PC-SPEC Licorice root
Bilberry Garlic St. John's Wart
Cayenne Ginger Yohimbe
Echinacea Gingko Biloba Vitamin E